

## Ozone therapy on Rats Submitted to Subtotal Nephrectomy: Role of Antioxidant System

José Luis Calunga,<sup>1</sup> Zullyt B. Zamora,<sup>1</sup> Aluet Borrego,<sup>1</sup> Sarahí del Río,<sup>2</sup> Ernesto Barber,<sup>3</sup>  
Silvia Menéndez,<sup>1</sup> Frank Hernández,<sup>1</sup> Teresita Montero,<sup>2</sup> and Dunia Taboada<sup>3</sup>

<sup>1</sup> Department of Biomedicine, Ozone Research Center, National Center for Scientific Research, PO Box 6414, Havana, Cuba ([ozono@infomed.sld.cu](mailto:ozono@infomed.sld.cu)).

<sup>2</sup> "Luis Díaz Soto" Military Medicine Institute, Havana City, Cuba.

<sup>3</sup> "Victoria de Girón" Institute of Basic and Preclinical Sciences, Havana City, Cuba

**Key words:** Ozone therapy, chronic renal failure, reactive oxygen species, antioxidant defense system.

### Abstract

Chronic renal failure (CRF) represents a world health problem. Ozone increases the endogenous antioxidant defense system, preserving the cell redox state. The aim of this study is to evaluate the effect of ozone/oxygen mixture in the renal function, morphology and biochemical parameters, in an experimental model of CRF (subtotal nephrectomy). Ozone/oxygen mixture was applied daily, by rectal insufflation (0.5 mg/kg) for 15 sessions after the nephrectomy. Renal function was evaluated, as well as different biochemical parameters, at the beginning and at the end of the study (10 weeks). Renal plasmatic flow (RPF), glomerular filtration rate (GFR), the urine excretion index, the sodium and potassium excretions (as a measured of tubular function) in the ozone group were similar to sham group. Nevertheless, nephrectomized rats without ozone (positive control group) showed the lowest RPF, GFR and urine excretion figures, as well as tubular function. Animals treated with ozone presented systolic arterial pressure (SAP) figures lower than the positive control group, but higher values with respect to sham group. Serum creatinine values and protein excretion in 24 hours in the ozone group were decreased comparing with nephrectomized rats, but still higher than normal values. Histological study demonstrated that animals treated with ozone presented less number of lesions in comparison with nephrectomized rats. Thiobarbituric acid reactive substances were significantly increased in nephrectomized and ozone-treated nephrectomized rats in comparison with sham group. In the positive control group, superoxide dismutase (SOD) and catalase (CAT) showed the lowest figures in comparison with the other groups. However, ozone/oxygen mixture induced a significant stimulation in the enzymatic activity of CAT, SOD and glutathione peroxidase, as well as reduced glutathione in relation with sham and positive control groups. In this animal model of CRF, ozone rectal administrations produced a delay in the advance of the disease, protecting the kidneys against vascular, hemorheological and oxidative mechanisms. This behavior suggests ozone therapy has a protective effect on renal tissue by down regulation of the oxidative stress present in CRF.

## Introduction

CRF represents a world health problem; once established, goes irreversibly to a final stage, provoking the patient death. In contrast with the capacity of the kidneys to regain function following acute renal injury, renal injury of a more prolonged nature often leads to progressive and irreversible destruction of nephron mass [1]. Such reduction of renal mass, in turn, causes structural and functional hypertrophy of surviving nephrons. This compensatory hypertrophy is due to an adaptive hyperfiltration mediated by increase in glomerular capillary pressures and flows. Eventually, these adaptations prove maladaptive, predisposing to sclerosis of the residual glomerular population [1-4]. The intrarenal vasculature is the most affected structure, preventing an appropriate blood flow, favoring the glomerular sclerosis [1-5]. For that reason, the improvement of the rheological properties of the blood could delay the progression of the CRF. Glomerulonephritis was the most common initiating cause of CRF in the past, possibly because of its more aggressive treatment. Now, diabetes mellitus and hypertensive renal diseases are leading causes of CRF. The inexorable course to renal failure often is accompanied by anemia, malnutrition, impaired metabolism of carbohydrates, fats and proteins, impaired platelet function and defective utilization of energy [1].

Reactive oxygen species (ROS) play a key intermediary role in the pathophysiologic processes of a wide variety of clinical and experimental renal diseases. It ranges from acute to chronic injuries, making the kidney as the site in which several unrelated diseases involves ROS [6]. ROS have been demonstrated to be capable of degrading glomerular basement membrane and inducing glomerular injury, characterized by impaired glomerular filtration and sieving function [7,8]. In order to eliminate toxic ROS, cells are equipped with various antioxidant defense systems. Therefore, the development of tissue injury depends upon the balance between ROS generation and tissue antioxidant defense mechanism [9].

Among various antioxidant systems equipped within aerobic cells, three antioxidant enzymes, superoxide dismutase (SOD), glutathione peroxidase (GSH-Px) and catalase (CAT) are major mechanisms to reduce local levels of ROS. Thus, these enzymes distributed in cytosol and/or mitochondria can abase primary ROS, such as superoxide anion (by SOD) and hydrogen peroxide (by GSH-Px and CAT) before they can interact to form more reactive cytotoxic metabolites (hydroxyl radical or hypochlorous acid, among others). Studies in the past demonstrated that glomerular antioxidant enzymes levels are modulated. Thus, the glomerular antioxidant enzymes are suggested to play an important role in the functional derangement induced by the ROS [10]. CRF is associated with depressed SOD and elevated NADPH oxidase expression, which can contribute to oxidative stress by increasing superoxide anion [11].

Another metabolic disturbance associated with CRF is hyperlipidemia, closely related with decreased removal and increase of triacylglycerol production. Up regulation of fatty acid synthase (FAS) gene expression reveals another factor involved in disturbed lipid metabolism in CRF. It seems that elevated plasma insulin and cytokine concentration could play an important role in the mechanism responsible for the increased FAS gene expression in CRF [12].

Taking into account some of the ozone therapeutic properties, such as, antiplatelet activity [13], enhancement of cell energy [14] and the increase of the antioxidant defense system [15-25], the aim of this paper is to evaluate the effect of ozone therapy in the renal function, morphology and biochemical parameters that measure oxidative stress in an experimental model of CRF.

## Materials and Methods

### Chemicals

All reagents used in determinations of GSH, SOD, CAT, GSH-Px and thiobarbituric acid reactive substances (TBARS) were purchased from Sigma Chemicals (St Louis, MO). Other reagents of analytical grade were obtained from normal commercial sources.

### Animals

Thirty young female Wistar rats (180-200 g) were maintained in an air filtered and temperature conditioned room (20-22 °C) with a relative humidity of 50-52 %. Rats were fed with standard laboratory chow and water *ad libitum* and were kept under an artificial light/dark cycle of 12 h. The experiments were performed in accordance with the ethical guidelines for investigations in laboratory animals and were approved by the Ethical Committee for Animal Experimentation of the National Center for Scientific Research, Havana, Cuba.

### Treatment Schedule and Surgical Procedure

Ozone (O<sub>3</sub>) was generated by OZOMED 01 equipment manufactured by the Ozone Research Center (Cuba). Ozone was obtained from medical grade oxygen by means of a silent electric discharge, representing about 3 % of the gas mixture (ozone/oxygen). The ozone concentration was measured by using an UV spectrophotometer at 254 nm. The ozone dose is the product of the ozone concentration, expressed as mg/L, by the gas mixture volume (L). By knowing the body weight of the rat, the ozone dose is calculated as 0.5 mg/kg.

Surgery was performed as previously described [26]. Ventral laparotomy was performed under aseptic conditions after anesthesia (sodium pentobarbital, 30 mg/kg intraperitoneal route). The right kidney was then removed, while two-thirds of the left kidney underwent acute infarction by ligation of two first-order branches of the main renal artery. Recovery from anesthesia and from the surgical procedure was complete within 24 h.

Animals were allocated randomly to 3 experimental groups of 10 animals each: 1-Sham group (negative control group), rats underwent a ventral laparotomy under anesthesia as described above. However, only handling of the renal pedicle without the removal of renal mass was performed. 2- Positive control group, rats were subjected to 5/6 renal ablation as described above. 3- Ozone group, rats were handled as in group 2 but also received, after the damage, 15 sessions of the gas mixture, composed of oxygen (O<sub>2</sub>) + O<sub>3</sub> (2.5-2.6 ml at a concentration of 50 mg/L, representing a dose of 0.5 mg/kg weight), by rectal insufflation, once per day.

### Sample Preparation

A day before the subtotal nephrectomy, all animals were housed in metabolic cages during 24 h, without food and water *ad libitum*. In all animals, the weight and the systolic arterial pressure (SAP) in the tail, as well as protein excretion were measured. All these procedures and measurements

were repeated, after the partial nephrectomy, once a week, during 10 weeks, time during which, the CRF continued its evolution. The time of the study was not prolonged for more than 10 weeks, avoiding the unpredictable death due to the final stage of the CRF. In the last day of evolution, plasmatic clearance of p-amino-hippurate (PAH) and inulin, in order to know the renal plasma flow (RPF) and the glomerular filtration rate (GFR), respectively, were determined using the method of unique injection (no urine) and the multicompartmental analysis of the plasmatic concentration curves in 9 blood samples [27]. Also, the urine excretion index (urine volume/water ingested), sodium and potassium excretions were determined in order to know tubular work. Creatinine in serum was determined in the final blood sample obtained by intracardiac puncture (2 ml of blood were extracted). Thereafter the animals were euthanized by ether anesthesia. The kidneys were dissected and immediately frozen at -20°C until analysis could be completed.

## Biochemical assays

PAH and inulin were determined in deproteinated plasma samples by cadmium sulfate, using for PAH a photocolometric technique, modified by Smith and Tinkelstein [28]. Inulin was measured by the direct method of resorcinol without alkaline treatment [29], corresponding calibration curves were used [30]. Proteins were calculated by the Biuret photocolometric technique using a Spectrophotometer (Shimadzu) [31]. Potassium and sodium urine concentrations were measured for the calculation of the excretions of both substances in a Carning flame photometer (model 400), using the method described by Oser [32]. Creatinine in plasma was measured in deproteinated filtrates by the method of sodium tungstate, using for its evaluation the method of picric acid modified by Brot [33].

Kidney homogenates were obtained using a tissue homogenator Ultraturrax T25 Polytron at 4°C. The homogenates were prepared by using a 100 mM KCl buffer (pH 7) containing EDTA 0.3 mM (1:10 w/v) for GSH, TBARS, GSH-Px and SOD determinations (buffer 1). The homogenates were spun down with a centrifuge at 600g for 60 min at 4°C. The supernatants were taken for the biochemical determinations. GSH was determined by a slightly modified version of Beutler method [34], using a spectrophotometer. One mL of the kidney homogenate, as described before, was mixed with 1.5 mL of 5 % metaphosphoric acid and centrifuged at 3000g for 10 min at room temperature. Fifty hundred  $\mu$ L of this acidic supernatant was mixed with 2 mL of 0.2 M phosphate buffer and 0.25 mL of 0.04 % 5,5'-Dithio-bis(-2-nitrobenzoic acid). Absorbance of the yellow solution was measured at 412 nm within 10 min. A molar extinction coefficient of  $13.6 \text{ M cm}^{-1}$  that describes the formation of the thiolate anion by the reaction of sulfhydryl groups with 5, 5'-Dithio-bis-2-nitrobenzoic acid (DTNB) at 412 nm was used to quantify GSH. Enzymatic activity of SOD was determined by a modified version of Minami and Yoshikawa method [35]. One unit of SOD enzymatic activity is equal to the amount of enzyme that diminishes the initial absorbance of nitro blue tetrazolium by 50 %. CAT was determined according to Evans and Diplock method [36]. Kidney homogenates for CAT enzymatic assay were carried out using 50 mM phosphate buffer (pH 7) containing 1% Triton X-100 (1:9 w/v) (buffer 2). The enzyme activity is expressed as the first order constant that describes the decomposition of  $\text{H}_2\text{O}_2$  at room temperature. Enzymatic activity of GSH-Px was measured using a modified version of Thonson et al method [37]. The enzyme activity is expressed as international units of enzymatic activity/mg of protein. International units are expressed as  $\mu$ moles of hydroperoxides transformed/min/mL of enzyme. To estimate TBARS levels, a method described by Ohkawa et al [38] was used. The absorbance of 3 mL of the colored layer was measured spectrophotometrically at 532 nm, using 1,1,3,3-tetraethoxypropane as standard. Protein concentrations were determined by the method of Lowry [39], using bovine serum albumin as standard.

## Histological study

Samples of rat kidneys were taken and fixed in 10 % neutral buffered formalin, processed and embedded in paraffin. A pathologist unaware of the treatment schedule examined the histological sections, stained with hematoxylin and eosin.

## Statistical analysis

First, the OUTLIERS preliminary tests for detection of error values was used. Afterward, the one way analysis of variance (ANOVA), followed by homogeneity variance test (Bartlett-Box) were applied. In addition, Duncan's multiple range test and the Student's t test, for the comparison of two groups, were done. Results are presented as mean  $\pm$  standard deviation (SD). Different letters indicate a statistical significance of at least  $p < 0.05$ .

## Results and Discussion

At the end of the study (10 weeks after the partial nephrectomy), animals treated with ozone presented: SAP figures lower than the positive control group, but higher values with respect to sham group (negative control group). Urine excretion index, in the ozone group, was similar to the negative control group, however, positive control group presented lower values respect to the others groups (Table 1). Respect to protein excretion and serum creatinine figures, in the ozone group, were higher than the negative control group, but lower than the positive control group. Potassium and sodium excretion values, in negative control and ozone groups were similar, but lower in the positive group with respect to negative and ozone groups (Table 1). RPF and GFR in ozone and sham groups presented similar figures, but they are higher respect to positive control group (Table 1).

*Table 1. Behavior of the systolic arterial pressure (SAP); urine excretion index; protein, potassium and sodium excretions; creatinine figures; renal plasmatic flow (RPF) and glomerular filtration rate (GFR), at the end of the study, in the different groups.*

Measurements	Groups		
	Sham	Positive control	Ozone
SAP (mmHg)	111 $\pm$ 13 <sup>a</sup>	165 $\pm$ 13 <sup>b</sup>	130 $\pm$ 13 <sup>b</sup>
Urine excretion index	1.26 $\pm$ 0.20 <sup>a</sup>	0.82 $\pm$ 0.14 <sup>b</sup>	1.20 $\pm$ 0.22 <sup>a</sup>
Protein excretion (mg/24h/100g rat)	1.31 $\pm$ 0.23 <sup>a</sup>	8.78 $\pm$ 2.06 <sup>b</sup>	4.08 $\pm$ 0.85 <sup>c</sup>
Potassium exc. (mEq/24h/100g rat)	34.26 $\pm$ 10.50 <sup>a</sup>	26.17 $\pm$ 8.30 <sup>b</sup>	30.55 $\pm$ 9.25 <sup>a</sup>
Sodium exc. (mEq/24h/100g rat)	575 $\pm$ 55 <sup>a</sup>	440 $\pm$ 60 <sup>b</sup>	507 $\pm$ 40 <sup>a</sup>
Plasma creatinine ( $\mu$ mol/l)	95 $\pm$ 15 <sup>a</sup>	160 $\pm$ 11 <sup>b</sup>	106 $\pm$ 10 <sup>c</sup>
RPF (ml/min/100g-rat)	1.71 $\pm$ 0.25 <sup>a</sup>	1.22 $\pm$ 0.67 <sup>b</sup>	1.85 $\pm$ 0.22 <sup>a</sup>
GFR (ml/min/100g-rat)	0.34 $\pm$ 0.01 <sup>a</sup>	0.26 $\pm$ 0.06 <sup>b</sup>	0.36 $\pm$ 0.02 <sup>a</sup>

Data are mean  $\pm$  SD. Statistical significance among different letters of at least  $p < 0.05$ .

Histological renal injuries (RI) were 0, 100 and 13 % in the sham, positive control and ozone groups, respectively (Table 2). The histological findings for positive control and ozone groups were glomerular collapse (GC), tubule degeneration (TD) and cortical-medullar hemorrhages (CMH), with 100 % for the positive control group compared with 20, 10 and 10 %, respectively, for the ozone group. Convoluted tubules dilatation (CTD) and glomerular capsule dilatation (GCD) were similar for these groups, presented in 100 % of animals.


*Table 2. Histological findings in the residual renal mass, due to the partial nephrectomy, in the different experimental groups.*

Groups	GCD %	GC %	CTD %	TD %	CMH %	RI %
Sham	0	0	0	0	0	0
Positive control	100	100	100	100	100	100
Ozone	100	20*	100	10*	10*	13*

GCD-glomerular capsule dilatation; GC-glomerular collapse; CTD-convoluted tubules dilatation; TD-tubule degeneration; CMH-cortical-medullar hemorrhages; RI-renal injury. \* Statistical significance of at least  $p < 0.05$ .

The histological study of renal cortex is observed in figure 1 (a,b,c). GCD and CTD are reversible structural alterations that appear in the first stages of the renal damage by vascular ablation. However, GC, TD and CMH are lesions that have a significant effect in the glomerular-tubular relation, causing an irreversible damage in the renal function.

*Figure 1. Histological study of the renal cortex. A- Sham group (200X), normal morphology; B- Positive control group (200X), glomerular collapse, cortical hemorrhages, vascular congestion, dilatation of convoluted tubules and glomerular capsule dilatation; C- Ozone group (400X), a discreet dilatation of convoluted tubules.*



A

B

C

Subtotal nephrectomy induced a significant increase in TBARS ( $p=0.0253$ ) of 144 %, whereas applications of ozone/oxygen gaseous mixture after subtotal nephrectomy increase TBARS in 90 % ( $p=0.0253$ ) over the levels of positive control group (Table 3). SOD activity was significantly decreased in 55 % ( $p=0.0143$ ) in positive control group, but in ozone group, 39 % of the enzymatic activity was recovered ( $p=0.0253$ ). Ozone therapy after subtotal nephrectomy induced a total increase in SOD activity of 94 % ( $p=0.0281$ ). A similar behavior was observed for CAT enzymatic activity. In positive control group there was a significant decrease of 50 %, whereas in ozone group was observed a recovery of 218 % ( $p=0.0143$ ) in the enzymatic activity of CAT, indicating a total increase in CAT activity of 268 % ( $p=0.0034$ ). GSH concentration and GSH-Px enzymatic activity were not significantly affected by subtotal nephrectomy in this animal model, but when fifteen intrarectal applications of ozone/oxygen gaseous mixture were applied we observed a significant stimulation of GSH-Px, as well as an increase in GSH concentration. Renal GSH concentration was significantly increased in 87 % ( $p=0.0253$ ) whereas GSH-Px activity was increased in 39 % ( $p=0.0253$ ) (Table 3).

*Table 3. Renal concentration of different biochemical parameters at the end of the study in the experimental groups.*

Biochemical Parameters	Groups		
	Sham	Positive control	Ozone
TBARS (nmol/mg protein)	0.25 ± 0.03 <sup>a</sup>	0.61 ± 0.25 <sup>b</sup>	1.16 ± 0.32 <sup>c</sup>
CAT ( $k_{15}$ /g of wet tissue)	6.80 ± 0.39 <sup>a</sup>	3.40 ± 0.41 <sup>b</sup>	11.12 ± 2.47 <sup>c</sup>
SOD (units/mg protein)	8.72 ± 1.15 <sup>a</sup>	4.74 ± 0.22 <sup>b</sup>	16.60 ± 0.29 <sup>c</sup>
GSH(nmol/mg protein)	7.87 ± 0.90 <sup>a</sup>	7.84 ± 0.47 <sup>a</sup>	14.29 ± 1.20 <sup>b</sup>
GPx (UI/mg protein)	6.01 ± 0.25 <sup>a</sup>	7.53 ± 0.94 <sup>a</sup>	10.47 ± 1.22 <sup>b</sup>

TBARS- Thiobarbituric acid reactive substances; CAT- catalase; SOD- superoxide dismutase; GSH-reduced glutathione; GPx-glutathione peroxidase. One unit of SOD enzymatic activity is equal to the amount of enzyme that diminishes the initial absorbance of nitro blue tetrazolium by 50 % and CAT activity is described as the enzymatic activity quantity that transforms 1 mol of  $H_2O_2$ , at room temperature, at 15 min/g of wet tissue. International units are expressed as  $\mu$ moles of transformed hydroperoxides/min/mL of GPx. Statistical significance among different letters of at least  $p<0.05$ .

Animals submitted to the subtraction of 5/6 of the total renal mass moved forward the installation of the CRF, demonstrated by the decrease of RPF and GFR and the increase of SAP, plasma creatinine, protein excretion, decrease of potassium and sodium excretions, as well as the presence of renal damage in the histological study. This behavior is still more pronounced in the positive control group, where the renal damage achieved 100 %.

The results have shown, at the end of the study, that the animals treated with ozone had the highest figures of RPF and GFR, as well as lower figures of proteinuria, plasma creatinine concentration, higher urine excretion and lower SAP in comparison with the positive control group. These results can be linked to the ozone antiplatelet activity [10], diminishing blood viscosity, that could produced a decrease in the friction between the blood and the glomerular vascular walls, decreasing the flow resistance, increasing the RPF and GFR, achieving similar figures to those present in the sham group. The flow rise contributes to diminish the endothelial injuries and the glomerular collapse, avoiding the tubular hypoxia, the hemorrhages and the release of several proinflammatory cytokines [40-43].

A depression in SOD expression was reported previously [11] for the same CRF model that we used, that could be correlated with the observed diminution in the activity of SOD and CAT, which can contribute to oxidative stress by increasing superoxide anion and hydrogen peroxide generation.

As it was reported before [14-24), ozone therapy is an oxidative approach that provoking brief moments of oxidative stress may stimulate antioxidant system to fight against the phenomenon per se. Thus, the significant increase in lipid peroxidation, measured in the form of TBARS, was expected after the surgical procedure, followed by fifteen ozone applications. In spite of the significant increase in TBARS, an increase in antioxidant status was seen in the ozone group. The increase in renal TBARS might not be related with an increase in renal damage because the remarkable increased in antioxidant system surpass this effect, inducing a general status of antioxidant protection.

Renal damage in this model of CRF might correspond with generation of ROS species, such as superoxide anion and hydrogen peroxide. Renal GSH concentration and GSH-Px activity might not be directly affected by subtotal nephrectomy in this model. However, ozone therapy induced a significant increase, in renal GSH amount and GSH-Px activity ( $p=0.0253$ ), besides a remarkable stimulation of SOD and CAT activities, that surpass the increase in lipid peroxidation. This suggests that ozone therapy has a protective effect on renal tissue, by up regulation of the antioxidant system, protecting against the oxidative stress provoked by ischemia.

In the other hand, it had been demonstrated that ozone is able to regulate the calcium levels, maintaining its homeostasis, avoiding any damage to the cell structure [44]. Also, it is possible that the ozone therapy effect, with the stimulation of the antioxidant defense system [14-25], protected the tissues against the oxidative stress present in CRF [6], being in correspondence with the positive histological results obtained, where the ozone group showed the lowest percent of glomerular collapse, tubule degeneration and cortical-medullar hemorrhages compared with positive control group.

### Conclusion

In this animal model of CRF, rectal administrations of ozone produced a delay in the advance of the disease, protecting the kidneys against the deleterious effects present in the CRF. Consequently, whenever possible, ozone therapy may become an important therapy to improve the quality of life of patients suffering of CRF.

### References

1. Fauci AS, Braunwald E, Isselbacher KJ et al. Chronic Renal Failure. In: Braunwald E, Hauser SL, Fauci AS, Longo DL, Kasper DL, Jameson JL, eds. *Harrison's Principles of Internal Medicine*. New York, USA: McGraw-Hill Companies Inc., p.1513-1518, 1998.
2. Lane PH. Long term furosemide treatment in normal rat: dissociation of glomerular hypertrophy and glomerulosclerosis. *Am J Kid*. 33(6):1058-1065, 1999.
3. Fine LG. Adaptation of renal tubule in uremia. *Kidney Int*. 22:546-552, 1982.
4. Slomowitz L. Tubuloglomerular feedback in chronic renal failure. *Nephron* 45(4):264-269, 1987.
5. Sheldin DW, Gusbech G. The kidney physiology and pathophysiology. In: Roven Press, eds *The kidney physiology and pathophysiology*. New York, USA: Roven Press, p.1901-1905, 1985.
6. Ichikawa I., Kiyama S, Yoshioka T. Renal antioxidant enzymes: Their regulation and function. *Kidney Int*. 45:19-22, 1994.
7. Yoshioka T, Ichikawa I. Glomerular dysfunction induced by polymorphonuclear leukocyte-derived reactive oxygen species. *Am J Physiol*. 257(Renal Fluid Electrol. Physiol. 26):F53-F59, 1989.
8. Yoshioka T, Moore-Jarret T, Ichikawa I, Yared A. Reactive oxygen species of extra-renal origin can induce massive functional proteinuria, *Kidney Int*. 37:497-502, 1990.
9. Kawamura T, Yoshioka T, Bills T, Fogo A, Ichikawa I. Glucocorticoid activates glomerular antioxidant enzymes and protects glomeruli from oxidant injuries. *Kidney Int*. 40:291-301, 1991.
10. Yoshioka T, Bills T, Moore-Jarrett T, Greene HL, Burr IM, Ichikawa I. Role of antioxidant enzymes in renal oxidant injury. *Kidney Int*. 38:282-287, 1990.
11. Vaziri ND, Dicus M, Ho ND, Boroujerdi-Rad L, Sindhu RK. Oxidative stress and dysregulation of



- superoxide dismutase and NADPH oxidase in renal insufficiency. *Kidney Int.* 63(1):179-184, 2003.
12. Szolkiewicz M, Nieweglowski T, Korczynska J et al. Upregulation of fatty acid synthase gene expression in experimental chronic renal failure. *Metabolism* 51(12):1605-1610, 2002.
  13. Matsuno K, Miura T, Shinriki N. The effect of ozone on platelet activation, Proceedings of the 13th Ozone World Congress (Kyoto, Japan); 3:178-184, 1997.
  14. Bocci V. Ozone as a bioregulator. Pharmacology and toxicology of ozonotherapy today. *J Biol Regul Homeost Agents* 10:31-53, 1996.
  15. Barber E, Menéndez S, León OS et al. Prevention of renal injury after induction of ozone tolerance in rats submitted to warm ischemia. *Med Inflamm.* 8:37-41, 1999.
  16. Hernández F, Menéndez S, Wong R. Decrease of blood cholesterol and stimulation of antioxidative response in cardiopathy patients treated with endovenous ozone therapy. *Free Rad Biol Med.* 19:115-119, 1995.
  17. León OS, Menéndez S, Merino N et al. Ozone oxidative preconditioning: a protection against cellular damage by free radicals. *Med Inflamm.* 7:289-294, 1998.
  18. Peralta C, León OS, Xaus C et al. Protective effect of ozone treatment on the injury associated with hepatic ischemia-reperfusion: antioxidant-prooxidant balance. *Free Rad Res.* 31:191-196, 1999.
  19. Candelario-Jalil E, Mohammed-Al-Dalain S, León OS, Menéndez S, Pérez-Davidson G, Merino N. Oxidative preconditioning affords protection against carbon tetrachloride-induced glycogen depletion and oxidative stress in rats. *J Appl Toxicol.* 21:297-301, 2001.
  20. Ajamieh H, Merino N, Candelario-Jalil E et al. Similar protective effect of ischemic and ozone oxidative preconditionings in liver ischaemia/reperfusion injury. *Pharmacological Research* 45(4):333-339, 2002.
  21. Al-Dalain SM, Martínez G, Candelario-Jalil E, Menéndez S, León OS. Ozone treatment reduces markers of oxidative and endothelial damage in an experimental diabetes model in rats, *Pharmaceutical Research* 44(5):391-396, 2001.
  22. Ajamieh H H, Menéndez S, Martínez-Sánchez G et al. Effects of ozone oxidative preconditioning on nitric oxide generation and cellular redox balance in a rat model of hepatic ischaemia-reperfusion. *Liver Int.* 24:55-62, 2004.
  23. Ajamieh HH, Berlanga J, Merino N et al. Role of protein synthesis in the protection conferred by ozone oxidative preconditioning in hepatic ischemia/reperfusion. *Int Transplant* 18:604-612, 2005.
  24. Al-Dalain SM, Martínez G, Menéndez S et al. Therapeutic efficacy of ozone medical treatments in patients with diabetic foot. *European Journal of Pharmacology* 2005 (in press).
  25. Zamora Z, Borrego A, López O et al. Effects of ozone oxidative preconditioning on TNF- $\alpha$  release and antioxidant-prooxidant intracellular balance in mice during endotoxic shock. *Med Inflamm* 2005 (in press).
  26. Anderson S, Meyer TW, Rennke HG, Brenner BM. Control of glomerular hypertension limits glomerular injury in rats with reduced renal mass. *J Clin Invest.* 76:612-617, 1985.
  27. Folin O. A system of blood analysis. *J Biol Chem.* 38:81-85, 1949.
  28. Smith HW, Tinkelstein N. The renal clearance of substituted hipuric acid derivatives and other aromatic acids in dog and man. *J Clin Invest.* 24:388-393, 1945.
  29. Schreiner G. Determination of inulin by means of resorcinol. *Proc Soc Exper Biol and Med.* 70:726-730, 1950.
  30. Jamison RL, Massly RH. The uramic concentrating mechanism. *N Eng J Med.* 295:1059-1067, 1976.
  31. Oser BL. Determination of sodium and potassium. In: Hawk's, eds *Physiological Chemistry*. New York, USA : MacGraw-Hill, p. 1140-1145, 1965.
  32. Oser BL Biuret test for protein determination. In: Hawk's, eds *Physiological Chemistry*. New York, USA : MacGraw-Hill, p. 179-84, 1965.
  33. Brot J. The renal clearance endogenous creatinine in man. *J Clin Invest.* 7:64552, 1948.
  34. Beutler F, Duron O, Mikus B. Improved method for the determination of blood glutathione. *J Lab Clin Med.* 6:882-887, 1963.
  35. Minami M, Yoshikawa H. A simplified assay method of superoxide dismutase activity for clinical use. *Clin Chim Acta* 2:337-343, 1979.
  36. Evans C, Diplock AT. Techniques in Free Radical Research. In: Burtin RH and Knippenberg PH, eds *Laboratory Techniques in Biochemistry and Molecular Biology*. The Netherlands: Elsevier, p.199-206, 1991.
  37. Faraji B, Kang HK, Valentine JL. Methods compared for determining glutathione peroxidase activity in blood. *Clin Chem.* 33:539-544, 1987.
  38. Ohkawa H, Orishi N, Yagi K. Assay for lipid peroxidation in animals and tissues by thiobarbituric acid reaction. *Anal Biochem.* 95:351-7, 1979.
  39. Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. Protein measurement with the Folin phenol reagent. *J Biol Chem.* 193:165-169, 1951.
  40. Mezzano SA, Droguett M, Burgos ME. Overexpression of chemokines, fibrogenic cytokines, and myofibroblasts in human membranous nephropathy. *Kidney Int.* 57(1):147-151, 2000.
  41. Hansch GM, Wagner C, Burger A. Matrix protein synthesis by glomerular mesangial cells in culture: effects

- of transforming growth factor beta (TGF beta) and platelet-derived growth factor (PDGF) on collagen type IV mRNA. *J Cell Physiol.* 163(3):451-456, 1995.
42. Ito Y, Aten J, Bende RJ. Expression of connective tissue growth factor in human renal fibrosis. *Kidney Int.* 53(4):853-858, 1998.
  43. Fine LG, Orphanides C, Norman JT. Progressive renal disease: The chronic hypoxia hypothesis, *Kidney Int.* 53(Suppl.65):S74-S77, 1998.
  44. León OS, Menéndez S, Merino N et al. Influencia del precondicionamiento oxidativo con ozono sobre los niveles de calcio. *Revista CENIC Ciencias Biológicas* 29(3):134-136, 1998.